

Contemporary determinants of outcome after colorectal surgery

Citation for published version (APA):

Cakir, H. (2020). *Contemporary determinants of outcome after colorectal surgery: adherence in enhanced recovery and visceral obesity*. [Doctoral Thesis, Maastricht University]. Ridderprint BV.
<https://doi.org/10.26481/dis.20200110hc>

Document status and date:

Published: 01/01/2020

DOI:

[10.26481/dis.20200110hc](https://doi.org/10.26481/dis.20200110hc)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Valorisation

Valorisation

In this thesis we contributed evidence for the importance of adherence after implementation of an ERAS program. In short, the ERAS program is a multidisciplinary approach to reduce surgical stress leading to early recovery and shorten postoperative length of stay (LOS) in patients undergoing major surgery.¹ The ERAS program consists of evidence-based changes in preoperative, intra-operative and postoperative treatments.¹ Nowadays, the ERAS program is the standard care after colorectal surgery. Higher adherence to the ERAS program led to a reduced LOS and better outcome in elective colonic cancer surgery.²⁻⁴ Therefore, compliance with the ERAS elements is crucial for a shorter LOS and fewer complications in patients undergoing elective colon resections for malignancy.⁵ Despite the obvious benefits of the ERAS program, its implementation and adherence to the protocol remain a challenge. Adherence to an ERAS program is complex because of its multidisciplinary character; it is sensitive to change, and continuous monitoring and feedback are required to ensure ongoing success.⁵ In this era of increasing numbers of protocols and regulations, hospital administrations should take responsibility to ensure these multidisciplinary protocols are continuously implemented in an organization, because without adherence the care and cost benefits are easily lost.⁵ Key factors include repeated training sessions and the presence of trained nurse–practitioners at all stages in perioperative care.⁵ It is useful to continuously monitor barriers and facilitators for implementation and adherence.⁵ In addition, the nationwide introduction of the ERAS protocol has resulted in an unprecedented equalization of perioperative care for the colorectal cancer patient increasing the quality of multicenter scientific programs on surgical outcome

Another item affecting the outcome of colorectal surgery was visceral obesity. In short, obesity is an increasing health problem in the Western world. Visceral obesity has been associated with worse outcome in colorectal surgery and other forms of surgery.⁶⁻¹³ The measurement of visceral fat contributes to the risk assessment of postoperative problems in patients undergoing colonic surgery^{14,15} and should be investigated in a larger context where it is conceivable that the measurement becomes a standard part of the preoperative work-up for patients with a colonic malignancy. In the preoperative work-up of the colorectal cancer patient, the assessment of visceral fat could be performed easily with CT imaging of the abdomen.¹⁶ This information is also important in metabolic risk profiling the elective colorectal surgery patients and offers the opportunity to develop tailor-made intervention programs.¹⁶ In this thesis, the worse outcome seems even more present in patients with a BMI <25 kg/m² in combination

with visceral obesity undergoing colon cancer surgery and this group may deserve more preoperative attention.¹⁶

References

1. Kehlet H. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth* 1997;78(5):606-17. Review.
2. Ahmed J, Khan S, Lim M, Chandrasekaran TV, MacFie J. Enhanced recovery after surgery protocols – compliance and variations in practice during routine colorectal surgery. *Colorectal Dis* 2012;14: 1045-51.
3. Gustafsson UO, Hausel J, Thorell A, Ljungqvist O, Soop M, Nygren J. Adherence to the enhanced recovery after surgery protocol and outcomes after colorectal cancer surgery. *Arch Surg* 2011;146: 571-7.
4. Vlug MS, Wind J, Hollmann MW, Ubbink DT, Cense HA, Engel AF, et al. Laparoscopy in combination with fast track multimodal management is the best perioperative strategy in patients undergoing colonic surgery: a randomized clinical trial (LAFA-study). *Ann Surg* 2011; 254(6): 868–75.
5. Cakir H, van Stijn MF, Lopes Cardozo AM, Langenhorst BL, Schreurs WH, van der Ploeg TJ, et al. Adherence to Enhanced Recovery After Surgery and length of stay after colonic resection. *Colorectal Dis* 2013;15:1019-25.
6. Dai Z, Xu YC, Niu L. Obesity and colorectal cancer risk: a meta-analysis of cohort studies. *World J Gastroenterol* 2007;13 (31):4199-206.
7. Tsujinaka S, Konishi F, Kawamura YJ, Saito M, Tajima N, Tanaka O, Lefor AT. Visceral obesity predicts surgical outcomes after laparoscopic colectomy for sigmoid colon cancer. *Dis Colon Rectum* 2008; 51(12):1757–65.
8. Ueda J, Ichimiya H, Okido M, Kato M. The impact of visceral fat accumulation on laparoscopy-assisted distal gastrectomy for early gastric cancer. *J Laparoendosc Adv Surg Tech A* 2009;19(2):157–62.
9. Yoshikawa K, Shimada M, Kurita N, Iwata T, Nishioka M, Morimoto S, Miyatani T, Komatsu M, Mikami C, Kashiwara H. Visceral fat area is superior to body mass index as a predictive factor for risk with laparoscopy-assisted gastrectomy for gastric cancer. *Surg Endosc* 2011;25(12):3825–30.
10. Miyaki A, Imamura K, Kobayashi R, Takami M, Matsumoto J. Impact of visceral fat on laparoscopy-assisted distal gastrectomy. *Surgeon* 2013;11(2):76–81.
11. Tokunaga M, Hiki N, Fukunaga T, Ogura T, Miyata S, Yamaguchi T. Effect of individual fat areas on early surgical outcomes after open gastrectomy for gastric cancer. *Br J Surg* 2009;96(5):496–500.
12. Shimizu A, Tani M, Kawai M, Hirono S, Miyazawa M, Uchiyama K, Yamaue H (2011) Influence of visceral obesity for postoperative pulmonary complications after pancreaticoduodenectomy. *J Gastrointest Surg* 2011;15(8):1401–10.
13. Hagiwara M, Miyajima A, Hasegawa M, Jinzaki M, Kikuchi E, Nakagawa K, Oya M. Visceral obesity is a strong predictor of perioperative outcome in patients undergoing laparoscopic radical nephrectomy. *BJU Int* 2012;110(11): E980–E984
14. Ballian N, Lubner MG, Munoz A, Harms BA, Heise CP, Foley EF, Kennedy GD. Visceral obesity is associated with outcomes of total mesorectal excision for rectal adenocarcinoma. *J Surg Oncol* 2012;105(4):365–70.
15. Seki Y, Ohue M, Sekimoto M, Takiguchi S, Takemasa I, Ikeda M, Yamamoto H, Monden M. Evaluation of the technical difficulty performing laparoscopic resection of a rectosigmoid carcinoma: visceral fat reflects technical difficulty more accurately than body mass index. *Surg Endosc* 2007;21(6):929–34.
16. Cakir H, Heus C, Verduin WM, Lak A, Doodeman HJ, Bemelman WA, Houdijk AP. Visceral obesity, body mass index and risk of complications after colon cancer resection: A retrospective cohort study. *Surgery* 2015;157(5):909-15.